

Updates in Nutritional Disorders and Therapy

Editorial

Open Access

Dietary Interventions Reverse Insulin and Synaptic Plasticity Defects Linking to Diabetes and Neurodegenerative Diseases

Martins IJ*

¹Centre of Excellence in Alzheimer's Disease Research and Care, School of Medical Sciences, Edith Cowan University, Australia

²School of Psychiatry and Clinical Neurosciences, The University of Western Australia, Australia

³McCusker Alzheimer's Research Foundation, Holywood Medical Centre, Australia

***Corresponding author:** Dr. Ian J Martins, School of Medical Sciences, Edith Cowan University, 270 Joondalup Drive, Joondalup, Western Australia 6027, Australia, Tel: +61863042574; Email: i.martins@ecu.edu.au

Acknowledgments: This work was supported by grants from Edith Cowan University, the McCusker Alzheimer's Research Foundation, and the National Health and Medical Research Council.

Copyright: © 2017 Martins IJ. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source.

Original Submission

Received: March 07, 2017

Accepted: March 21, 2017

Published: April 07, 2017

Open Peer Review Status: Editorials, news items, analysis articles, and features do not undergo external peer review.

How to cite this article: Martins IJ. Dietary Interventions Reverse Insulin and Synaptic Plasticity Defects Linking to Diabetes and Neurodegenerative Diseases. Updates Nutr Disorders Ther. (2017) 1: 4.1

Updates in Nutritional Disorders and Therapy

Keywords

SNAP25; Insulin Secretion; Synaptic Plasticity; Diabetes; Lipopolysaccharides; Patulin

Editorial

In the developing world food and nutrition guidelines with relevance to nutrition and metabolism are required to reduce elevated cholesterol levels with relevance to diabetes, Parkinson's disease and Alzheimer's disease [1]. In diabetes and neurodegenerative diseases global non alcoholic fatty liver disease (NAFLD) has become of major concern with the reduced hepatic metabolism of toxic compounds that increase with age to milligrams quantities of bacterial lipopolysaccharides (LPS), mycotoxins and xenobiotics that induce neuron death. The increased brain content of these toxic compounds completely inactivate drug therapy with increased repetitive drug exposure to neurons and synapses [2-4]. In the global diabetes epidemic diabetes is expected to double by the year 2050 in the developing world and diets that reverse or stabilize mitochondrial apoptosis [5] have become relevant to synaptic plasticity and to prevent neuronal death [6]. These toxic compounds determine the survival of various species with relevance to greater toxicity to neurons and their synapses in diabetes when compared to toxic amyloid beta oligomers and alpha synuclein oligomers in Parkinson's disease and Alzheimer's disease [1].

In diabetes information processing is defective and related to defective interactions between neurons and synapses [7-10]. The toxic compounds that increase in the brain of diabetics induce post-translational modifications (Figure 1) between cholesterol interacting proteins [1] and are responsible for the poor synaptic connections with relevance to insulin resistance and impairments in synaptic plasticity. Defects in cholesterol transport in the brain are responsible defective information processing and synaptic plasticity in diabetes and neurodegenerative diseases. Interests in plasma LPS and patulin levels (Figure 1) in the developing world have attracted major concern with relevance to accelerated cross linking (intra and intermolecular crosslinks) between cholesterol interacting proteins [1] in cell membranes with relevance to cholesterol, alpha synuclein and amyloid beta metabolism (Figure 1). Cholesterol interacting proteins [1] include apolipoprotein AI (apo AI), apolipoprotein E (apo E), ATP binding cassette transport protein 1 (ABCA1), low density lipoprotein receptors (LDLr/LRP), G-coupled protein receptors (GPCR) (11), phospholipid transfer protein (PLTP), amyloid beta (A β) and alpha synuclein.




Figure 1: The poor survival rates of individuals in populations connected with increased global death rates has raised concerns for individuals in the developing world with relevance to the current global diabetes epidemic. Hypercholesterolemia is a major defect and related to post-translational modifications of cholesterol interacting proteins (cysteine residues). LPS/Patulin are responsible for inactivation of cysteine residues in cholesterol interacting proteins (apo AI, apo E, GPCR, PLTP, LDLr/LRP, SNARE/caveolin 1) with relevance to diabetes (SNAP25 (cysteine-rich)/insulin release), Parkinson's disease (SNAP 25/alpha synuclein) and Alzheimer's disease (SNAP25/amyloid beta).

Life changing scientific discoveries indicate that the nuclear receptor Sirtuin 1 (Sirt 1) is important to cholesterol homeostasis [12] and to prevent synaptic plasticity defects in diabetes, Parkinson's disease and Alzheimer's disease [7-10]. Post-transcriptional dysregulation (via cysteine residues) that involve zinc defective Sirt 1/p53 interactions [1,13-20] effect caveolin 1 expression [21] that is important to cellular cholesterol homeostasis, insulin receptor transport and activity [22-25]. Interference of synatosomal associated protein 25 (SNAP25)-caveolin 1 interaction corrupt neuron synapse interactions (Figure 1) with implication of food ingestion that contain (LPS/Patulin) that inactivate cysteine residues essential for and SNAP25 membrane interactions [25-27] and Sirt 1 function [12,13,28]. SNAP25 is involved in SNARE/caveolin 1 complex (Figure 1) with relevance to synaptic transmission in diabetes and neurodegenerative diseases [29-33]. Interference with SNAP25 and SNARE proteins in membranes now implicate SNAP25 to be involved with pancreatic beta cell insulin secretion (Figure 1) release with SNAP25a and SNAP25b now relevant to metabolic disease, synaptic plasticity and accelerated neurodegeneration [34-40].

Nutritional interventions in the developing world need to be implemented to prevent a miscible disease with relevance to diabetes, Parkinson's disease and Alzheimer's disease. LPS levels should be carefully monitored in various foods such as fruits, bread, apple/apple juice, cereals, cheese and nuts (Figure 2). Patulin levels should be assessed in yoghurt, cream, cheese, bread, fruits and unpasteurized fruit juices (Figure 2). In miscible diseases such as diabetes and neurodegenerative diseases Ochratoxin A levels should be assessed in coffee and alcohol to determine effects on neuron and mitochondria apoptosis. The plasma levels of LPS and patulin in the developing world (Figure 2) should be determined to prevent a miscible disease such as diabetes and neurodegeneration (synaptic plasticity defects). In the developing world low fat and very low carbohydrate diets are encouraged to prevent absorption of lipophilic LPS and

Updates in Nutritional Disorders and Therapy

patulin that are present in lipoproteins with toxic components transported to various tissues such as the pancreas (insulin secretin disorders), adipose tissue, liver (NAFLD) and brain (synaptic plasticity defects).

PATULIN	LPS	OCHRATOXIN A
FRESH/DRIED FRUITS BREAD (PROCESSING STAGES) APPLE/APPLE JUICE APPLE RIPENING CEREALS BLUE CHEESE CHEESE SPOILAGE NUT/NUT PRODUCTS	UNPASTEURIZED FRUIT JUICES CIDERS POORLY OVEN BAKED BREAD FRUITS FRENCH CHESSE YOGHURT CREAM VEGETABLES	COFFEE CHEESE (RIND/INNER) ALCOHOL (BEER/WINE)
<p>SCIENTIFIC WARNING: Patulin may have been used as an antibiotic against Gram-positive and Gram-negative bacteria infection</p> <p>Previous reports of its use warn of toxicity</p>		

Figure 2: Blood tests for LPS and patulin in diabetic individuals in the developing world are required to determine the diagnosis of a miscible disease with relevance to diabetes and neurodegeneration. Nutritional intervention is required to measure LPS, Patulin and ochratoxin A in various foods to prevent ingestion of the end products that induce the miscible disease with relevance to mitochondrial dysfunction (NAFLD) and neuron apoptosis. In the developing world Patulin should not be ingested as an antibiotic with relevance to its poisonous effects (NAFLD) in various individuals.

Conclusion

In the developing world the global diabetes epidemic is predicted to escalate over the next 30 years and a miscible disease with relevance to brain synaptic plasticity defects is expected to affect many of these diabetic individuals. Dietary interventions and nutritional research indicate that various food end products such as LPS and mycotoxins should not enter the blood plasma in diabetic individuals. LPS and patulin effect various cholesterol interacting proteins with effects on pancreatic insulin release and brain synaptic transmissions. Stabilization of this miscible disease with relevance to diabetes and neurodegeneration involves low calorie nutrition that activate nuclear receptors relevant to the metabolism of LPS/mycotoxins with therapeutic activation of brain drug therapy in diabetic individuals in the developing world.

References

1. Martins IJ. Overnutrition Determines LPS Regulation of Mycotoxin Induced Neurotoxicity in Neurodegenerative Diseases. *Int J Mol Sci.* 2015;16: 29554–29573.
2. Jones S, Bonci A. Synaptic plasticity and drug addiction. *Curr Opin Pharmacol.* 2005;5:20-25.
3. Grillo CA, Piroli GG, Lawrence RC, Wrighten SA, Green AJ, et al. Hippocampal Insulin Resistance Impairs Spatial Learning and Synaptic Plasticity. *Diabetes.* 2015;64:3927-3936.
4. Lüscher C, Malenka RC. Drug-evoked synaptic plas-

ticity in addiction: from molecular changes to circuit remodeling. *Neuron.* 2011;69:650-663.

5. Carvalho C, Santos MS, Oliveira CR, Moreira PI. Alzheimer's disease and type 2 diabetes-related alterations in brain mitochondria, autophagy and synaptic markers. *Biochim Biophys Acta.* 2015;1852:1665-1675.
6. Martins IJ. Diet and Nutrition reverse Type 3 Diabetes and Accelerated Aging linked to Global chronic diseases. *J Diab Res Ther.* 2016; 2: 1-6.
7. Stranahan AM, Norman ED, Lee K, Cutler RG, Telljohann RS, et al. Diet-induced insulin resistance impairs hippocampal synaptic plasticity and cognition in middle-aged rats. *Hippocampus.* 2008;18:1085-1088.
8. Gispen WH, Biessels GJ. Cognition and synaptic plasticity in diabetes mellitus. *Trends Neurosci.* 2000;23:542-549.
9. Biessels GJ, Kamal A, Ramakers GM, Urban IJ, Spruijt BM, et al. Place learning and hippocampal synaptic plasticity in streptozotocin-induced diabetic rats. *Diabetes.* 1996;45:1259-1266.
10. Colicos MA, Syed NI. Neuronal networks and synaptic plasticity: understanding complex system dynamics by interfacing neurons with silicon technologies. *J Exp Biol.* 2006;209:2312-2319.
11. Gimpl G. Interaction of G protein coupled receptors and cholesterol. *Chem Phys Lipids.* 2016;199:61-73.
12. Michán S, Li Y, Chou MM, Parrella E, Ge H, et al. SIRT1 is essential for normal cognitive function and synaptic plasticity. *J Neurosci.* 2010;30:9695-9707.
13. Ng F, Wijaya L, Tang BL. SIRT1 in the brain—connections with aging-associated disorders and lifespan. *Front Cell Neurosci.* 2015; 9:1-13.
14. Martins IJ. Nutritional and Genotoxic Stress Contributes to Diabetes and Neurodegenerative Diseases such as Parkinson's and Alzheimer's Diseases. *Frontiers in Clinical Drug Research -CNS and Neurological Disorders.* 2015;3:158-192.
15. Hainaut P, Mann K. Zinc binding and redox control of p53 structure and function. *Antioxid Redox Signal.* 2001;3:611-623.
16. Vergnano AM, Rebola N, Savtchenko LP, Pinheiro PS, Casado M, et al. Zinc dynamics and action at excitatory synapses. *Neuron.* 2014;82:1101-1114.
17. Pace NJ, Weerapana E. Zinc-Binding Cysteines: Diverse Functions and Structural Motifs. *Biomolecules.* 2014; 4: 419–434.
18. Rainwater R, Parks D, Anderson ME, Tegtmeyer P, Mann K. Role of cysteine residues in regulation of p53

Updates in Nutritional Disorders and Therapy

- function. *Mol Cell Biol.* 1995;15:3892-3903.
19. Kaar JL, Basse N, Joerger AC, Stephens E, Rutherford TJ, et al. Stabilization of mutant p53 via alkylation of cysteines and effects on DNA binding. *Protein Sci.* 2010;19:2267-2278.
 20. Stoner CS, Pearson GD, Koç A, Merwin JR, Lopez NI, et al. Effect of thioredoxin deletion and p53 cysteine replacement on human p53 activity in wild-type and thioredoxin reductase null yeast. *Biochemistry.* 2009;48:9156-9169.
 21. Frank PG, Cheung MW, Pavlides S, Llaverias G, Park DS, et al. Caveolin-1 and regulation of cellular cholesterol homeostasis. *Am J Physiol Heart Circ Physiol.* 2006;291:H677-686.
 22. Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in insulin signaling and diabetes. *Am J Physiol Endocrinol Metab.* 2003; 285:E1151–E1160.
 23. Yamamoto M, Toya Y, Schwencke C, Lisanti MP, Myers MG et al. Caveolin is an activator of insulin receptor signaling. *J. Biol. Chem.* 1998;273:26962–26968.
 24. Nystrom FH, Chen H, Cong LN, Li Y, Quon MJ. Caveolin-1 interacts with the insulin receptor and can differentially modulate insulin signaling in transfected Cos-7 cells and rat adipose cells. *Mol. Endocrinol.* 1999;13:2013–2024.
 25. Cohen AW, Razani B, Wang XB, Combs TP, Williams TM et al. Caveolin-1-deficient mice show insulin resistance and defective insulin receptor protein expression in adipose tissue. *Am. J. Physiol. Cell Physiol.* 2003; 285: C222–C235.
 26. Washbourne P, Cansino V, Mathews JR, Graham M, Burgoyne RD, et al. Cysteine residues of SNAP-25 are required for SNARE disassembly and exocytosis, but not for membrane targeting. *Biochem J.* 2001;357:625-634.
 27. Greaves J, Prescott GR, Fukata Y, Fukata M, Salaun C, et al. The hydrophobic cysteine-rich domain of SNAP25 couples with downstream residues to mediate membrane interactions and recognition by DHHC palmitoyl transferases. *Mol. Biol. Cell.* 2009;20:1845-1854.
 28. ZeeRS, Pimentel DR, Hou, XL, Zang M, Yao, CX, et al. Sirtuin-1 zinc thiolate center is a direct molecular target of oxidants. In *Proceedings of the Society for Free Radical Biology and Medicine's 15th Annual Meeting, Indianapolis, IN, USA, 19–23 November 2008. Free Radical Biology and Medicine.* 2008; 45: S76.
 29. Magga JM, Kay JG, Davy A, Poulton NP, Robbins SM, et al. ATP dependence of the SNARE/caveolin 1 interaction in the hippocampus. *Biochem Biophys Res Commun.* 2002;291:1232-1238.
 30. Braun JE, Madison DV. A novel SNAP25-caveolin complex correlates with the onset of persistent synaptic potentiation. *J Neurosci.* 2000;20:5997-6006.
 31. Yamamori S, Itakura M, Sugaya D, Katsumata O, Sakagami H, et al. Differential expression of SNAP-25 family proteins in the mouse brain. *J Comp Neurol.* 2011;519:916-932.
 32. Südhof TC, Rizo J. Synaptic Vesicle Exocytosis. *Cold Spring Harb Perspect Biol.* 2011; 3: a005637.
 33. Antonucci F, Corradini I, Fossati G, Tomasoni R, Menna E, et al. SNAP-25, a Known Presynaptic Protein with Emerging Postsynaptic Functions. *Front Synaptic Neurosci.* 2016; 8: 7.
 34. Gonelle-Gispert C, Molinete M, Halban PA, Sadoul K. Membrane localization and biological activity of SNAP-25 cysteine mutants in insulin-secreting cells. *J Cell Sci.* 2000;113:3197-3205.
 35. Nagamatsu S, Nakamichi Y, Yamamura C, Matsushima S, Watanabe T, et al. Decreased expression of t-SNARE, syntaxin 1, and SNAP-25 in pancreatic beta-cells is involved in impaired insulin secretion from diabetic GK rat islets: restoration of decreased t-SNARE proteins improves impaired insulin secretion. *Diabetes.* 1999;48:2367-2373.
 36. Valladolid-Acebes I, Daraio T, Brismar K, Harkany T, Ögren SO, et al. Replacing SNAP-25b with SNAP-25a expression results in metabolic disease. *Proc Natl Acad Sci U S A.* 2015; 112: E4326-4335.
 37. Chauhan NB, Siegel GJ. Reversal of amyloid beta toxicity in Alzheimer's disease model Tg2576 by intraventricular anti-amyloid beta antibody. *J Neurosci Res.* 2002;69:10-23.
 38. Brinkmalm A, Brinkmalm G, Honer WG, Frölich L, Hausner L, et al. SNAP-25 is a promising novel cerebrospinal fluid biomarker for synapse degeneration in Alzheimer's disease. *Mol Neurodegener.* 2014;9:53.
 39. Thumond DC. Regulation of Insulin Action and Insulin Secretion by SNARE-Mediated Vesicle Exocytosis. *Madame Curie Bioscience Database [Internet]. Austin: Landes Bioscience.* 2000-2013.
 40. Al-Daghri NM, Costa AS, Alokail MS, Zanzottera M, Alenad AM, et al. Synaptosomal Protein of 25 kDa (Snap25) Polymorphisms Associated with Glycemic Parameters in Type 2 Diabetes Patients. *Journal of Diabetes Research.* 2016; 2016: 8943092.